

NOTAS

Biomass production and carbon balance of a short rotation forestry of *Populus deltoides* (clone Lux) under two different cutting cycles

Producción de biomasa y balance de carbono en una rotación de *Populus deltoides* (clon Lux) bajo dos ciclos de corta

Antonio María Cabrera ^{}, Cristiano Tozzini ^b, Sergio Espinoza ^a, Rómulo Santelices ^a, Fernando Meza ^c**

^{*}Corresponding autor: ^aUniversidad Católica del Maule, Centro de Desarrollo del Secano Interior, Avenida San Miguel 3605, Talca, Chile, phone +56-071-2413728, acabrera@ucm.cl

^bCentro Interdipartimentale di Ricerche Agro-ambientali “Enrico Avanzi” via Vecchia di Marina, 6 - 56122 San Piero a Grado (Pisa), Italy.

^cUniversidad Santo Tomás, Facultad de Ciencias Básicas, Departamento de Ciencias Básicas, Mendoza # 120, Los Ángeles, Chile.

SUMMARY

The emissions of green house gases contribute to global warming. Consequently, renewable energies such as solar, wind power, hydropower and bio energy appear as alternatives for reducing these emissions. One promising source of renewable energy is forest biomass, for it is considered to have neutral CO₂ balance. The problem is to quantify the amount of CO₂ that is emitted for a certain amount of energy generated, compared to the CO₂ emitted to generate the same amount of energy from a non-renewable source. In this paper we calculate biomass production and carbon dioxide balance of an eight-year-old *Populus deltoides* short rotation forestry (SRF) under two cutting cycles (harvesting every two years, biennial and every three years, triennial). We also compare the amount of energy generated per kg of CO₂ emitted in the use of fossil fuels versus the amount of energy generated per kilogram of CO₂ emitted. On the one hand, total biomass yield was 12.6 ± 0.9 Mg ha⁻¹ year⁻¹ and 15.1 ± 1.5 Mg ha⁻¹ year⁻¹ for biennial and triennial cutting cycles respectively. On the other hand, total emissions of CO₂ for growing a poplar SRF with a cycle of eight year and biennial cutting sequence reached a value of 738.8 kg ha⁻¹ year⁻¹; however, triennial short total emissions are equivalent to 695.5 kg ha⁻¹ year⁻¹. We concluded that the use of biomass makes a positive contribution to the reduction of greenhouse gases, in particular CO₂, reducing emissions almost five times compared with fossil fuels.

Key words: carbon dioxide balance, *Populus*, short rotation forestry, biomass production.

RESUMEN

Las emisiones de gases de efecto invernadero contribuyen al calentamiento global. En esta situación, las energías renovables como la solar, eólica, hidroeléctrica y biomasa, aparecen como una alternativa para reducir estas emisiones. Una prometedora fuente de energía renovable es la biomasa, considerada con un balance de CO₂ neutro. El problema es cuantificar el CO₂ emitido para una cierta cantidad de energía generada, en comparación con el CO₂ emitido para generar la misma cantidad de energía de una fuente no renovable. En este trabajo se calculó la producción de biomasa y el balance de CO₂ en una rotación de ocho años de edad de *Populus deltoides* bajo dos ciclos de corte (bienal y trienal). También se comparó la cantidad de energía generada por kilogramo de CO₂ emitido en el uso de combustibles fósiles frente a la cantidad de energía generada por kilogramo de CO₂ emitido en el uso de biomasa como energía. El rendimiento total de la biomasa fue de 12,6 ± 0,9 Mg ha⁻¹ año⁻¹ y 15,1 ± 1,5 Mg ha⁻¹ año⁻¹ para los ciclos bienal y trienal respectivamente. Las emisiones totales de CO₂ fueron de 738,8 kg ha⁻¹ año⁻¹ y 695,5 kg ha⁻¹ año⁻¹ para los ciclos bienal y trienal, respectivamente. Se concluyó que el uso de biomasa constituye una contribución positiva a la reducción de gases de efecto invernadero, en particular CO₂, reduciendo las emisiones casi cinco veces en comparación con combustibles fósiles.

Palabras clave: balance de dióxido de carbono, *Populus*, ciclos cortos de rotación, producción de biomasa.

INTRODUCTION

During the last years the strong growth of population globally recorded has resulted in a growing need for ener-

gy in developed and developing countries (Tlili 2015). The main risk from the growing pressure exerted by mankind on the use of energy resources (*i.e.* oil, gas, coal) is represented not only because they are finite resources, but also

for the impact on the environment caused by the use of these resources (ENEA 1999). In fact, the use of fossil fuels, such as gas, produces carbon dioxide (CO₂); which, in certain concentrations, is considered as one of the main causes of the greenhouse effect, which is the basis of global warming and climate change (IPCC 2006). As the atmospheric concentration of CO₂ grows, there is an increasing interest in restraining this growth to minimize potential impacts on the global climate (West and Marland 2002). One of the most promising alternatives for reducing CO₂ emissions is the production of biomass energy, specifically lignocellulosic crops that can be used to produce heat and electricity through direct combustion, or biofuel or biogas production through pyrolysis and gasification (Mantineo *et al.* 2009). Biomass energy, both lignocellulosic and herbaceous, is presented as a sustainable and environmentally friendly source of an alternative energy; in fact the production of energy from renewable sources has a limited environmental impact. In particular, the contribution to the greenhouse effect is extremely small: CO₂ emissions related to the stages of conversion processes are almost completely balanced by the CO₂ consumption required for the growth of the biomass used as an energy source (Cannell 2002).

One of the main problems associated with the introduction of alternative crops for energy production is to assess the long-term sustainability of these crops. For this purpose, CO₂ balances, energy balance and water use efficiency, among others, may represent an optimal tool for sustainability assessment in particular for the energy crops cultivation system, which has to ensure a reduction in CO₂ emissions, compared to the use of fossil fuels, *i.e.*, positive CO₂ balance.

Regarding the above mentioned, CO₂ balance has been widely used (Lewandowski *et al.* 1995, Cannell 2002, Kaur 2002, Hoosbeek *et al.* 2006) and it estimates the emissions generated in the whole crop production cycle compared with the CO₂ fixed by the plant during its growth. CO₂ balance takes into account all the issues arising from the implementation of the different forestry operations; considering not only the emissions produced during the combustion of biomass, but also the emissions generated in input productions (fertilizers, herbicides, etc.). For example, Dornburg *et al.* (2005), for a poplar SRF held in Poland and the Netherlands with fertilization every four years, obtained CO₂ emission values, in the production phase of the crop, in a range of 1,170-1,660 kg ha⁻¹ year⁻¹. The results in this study demonstrate to be consistent with most of the available data, which indicates a strong relationship between the cutting cycle and the productivity of the stand (Deckmyn *et al.* 2004). The objective is to estimate the eventual reduction of CO₂ emissions through the energy production from biomass compared to the same amount of energy produced from fossil fuel.

This paper hypothesizes that the CO₂ balance in an eight-year-old *Populus deltoides* Marshall (clone Lux) short rotation forestry under two cutting cycles is negative, *i.e.* the CO₂ fixed by the plants is higher than the CO₂ emitted in the entire production process.

METHODS

Study area. The study was conducted at the Centro di Ricerche Agro-ambientali (CIRAA) Enrico Avanzi at Pisa University (Italy). The experimental field is situated in San Piero a Grado, 43°40' N, 10°21' E at 5 m above sea level and 5 km far from the sea. The soil was a Xerofluvent (clay 20.1 %, silt 40.5 %, sand 39.4 %), typical of the lower River Arno, which is an alluvial plain characterized by a superficial water table (1.8 m deep in the driest conditions) and good nutrient availability (organic matter 1.8 %, total nitrogen content 1.3 g kg⁻¹, available phosphorus 8.8 mg kg⁻¹ and exchangeable potassium 128.3 mg kg⁻¹).

The average climate conditions of the site during the trial are shown in figure 1. During the experimental period (from 2000 to 2008), a considerable variability in rainfall was observed from year to year with a mean annual rainfall of approximately 750 mm (from 655 to 936 mm).

Experimental setup. The experiment began in the year 2000 planting 10,000 plants per hectare; it was conducted in two plots of 5,000 m² each. In 2002 both plots were harvested and it was in that year when two cutting cycles were chosen (biennial T2, and triennial T3). Each cutting cycle was represented in an individual plot, being each plot the basis for future comparisons. T2 was harvested in 2002, 2004, 2006 and 2008 and T3 was harvested in 2002, 2005 and 2008 (Cabrera *et al.* 2014). The soil was prepared ahead the planting season (October 1999) by deep plowing (50 cm), followed by sub soiling with ridge and plow. Then a pre-planting herbicide (Pendimetalin Click® 50, 3L ha⁻¹) and a fertilization using 600 kg ha⁻¹ of a generic 8-24-24 fertilizer was applied. In the year 2000 we planted 20 cm long uprooted *P. deltoides* (clone Lux) cuttings. The plant density was equivalent to 10,000 plants ha⁻¹ and the planting design was 2m x 0.5m. In 2002, the first harvest was made in both plots. The biennial (T2) was harvested in 2002, 2004, 2006 and 2008 and the triennial (T3) in 2002, 2005 and 2008; always at the end of February before the vegetative regrowth. To harvest, a harvesting took-loader-chipper, as well as a trailer carried by a tractor for the collection of chipped material, was used. After each harvest, supplementary fertilization applying 100 kg ha⁻¹ of nitrogen supplied as urea was applied.

At the end of the experiment (*i.e.* year 2008) the soil was recovered by removing the stumps with stump cutter and making a deep plowing and sub soiling. We used various types of tractors depending on the operation to perform: 132 kW of power for plowing and harvesting, 73 kW to sub soiling and 48 kW for all the other operations.

Biomass production. At every harvest, the fresh above-ground biomass of the whole plant was assessed. Data were sampled in five transects (5m long, 10 plants) per plot, which were always located within the three central rows of the plot. Biomass samples of each plot were oven




Figure 1. Climate conditions at the field experimental station. T_{max}: maximum temperature; T_{min}: minimum temperature; T_{med}: medium temperature.

Condiciones climáticas en el sitio de experimentación. T_{max}: temperatura máxima, T_{min}: Temperatura mínima; T_{med}: temperatura media.

dried at 105 °C (to constant mass) and the dry matter content was determined. Total biomass was calculated by multiplying the average dry weight of the plants of each transects by the plantation density. Average biomass per hectare for each cutting cycle was calculated by averaging the biomass yield in each harvest by the eight years of the trail duration. For dry weight, a Student's t-test was used to compare means under different treatments.

CO₂ balance. This study considered CO₂ emissions relating to the operations required to grow poplar short rotation forestry and turn harvested biomass into usable solid biofuel (woody chip) in the field. Post processing and the conversion of woodchips were excluded from the analysis. The propagation material CO₂ emissions were considered negligible.

The calculation of CO₂ balance begins with the calculation of the emissions for the different operations made to the soil and to the crop, starting with fuel and lubricant that characterizes each of these operations. Specific consumption for each of the farming operations were obtained from Bonari *et al.* (1999), for the same types of operations (table 1) in the same areas. Regarding harvesting, the data have been determined directly during the development of the experiment. The CO₂ emissions generated by the inputs (fuel, lubricants, fertilizers, herbicides, etc.) were calculated using coefficients found in literature (table 2).

To compare the CO₂ emissions generated with the potential energy obtainable from biomass, the energy output was determined subsequently by multiplying the dry mat-

ter yield by the calorific value of the plant material that is considered 18 MJ kg⁻¹ according to Cabrera (2009) and it is in the range of the most recent literature (Carmona *et al.* 2015). Thus, we are able to calculate the quantity of energy for every CO₂ kg emitted.

Table 1. Fuel and oil consumption of different agricultural operations and its related CO₂ emissions (Bonari *et al.* 1999).

Consumo de diesel, aceite y emisiones de CO₂ de diferentes operaciones agrícolas (Bonari *et al.* 1999).

Operation	Consumption			
	Diesel (kg ha ⁻¹)	Lubricant (kg ha ⁻¹)	CO ₂ emissions (kg ha ⁻¹)	
Deep plowing (50 cm)	65.0	0.36	207.2	
Subsoiling	14.6	0.04	46.5	
Plow	50.0	0.20	180.4	
Herbicide	7.0	0.05	22.3	
Fertilize	7.8	0.02	24.9	
Planting	18.7	0.04	59.6	
Harvest	T2	51.0	0.35	162.6
	T3	80.0	0.45	255
Stump cutter	6.2	0.03	19.8	
Subsoiling	14.6	0.04	46.4	

Table 2. Conversion factors for CO₂ emissions of different inputs.
 Factores de conversión en CO₂ de diferentes *inputs*.

Inputs	Conversion factor (kg kg ⁻¹)	Reference
Fuel	3.19	IPCC (2006)
Lubricant	2.95	IPCC (2006)
N	3.15	West and Marland (2002)
P ₂ O ₅	6.04	West and Marland (2002)
K ₂ O	4.39	West and Marland (2002)
Herbicide	15.92	West and Marland (2002)

RESULTS

Biomass production. The annual average production in the study of the complete cycle of *P. deltoides* clone Lux differs among treatments, being T3 the most productive cutting cycle (figure 2 and table 3).

Lower yields in the period 2004-2006 may be because rainfall in that period was lower than the average rainfall.

CO₂ balance. The calculation of CO₂ emissions for the poplar cultivation with biennial and triennial cutting cycles has been made considering all the emissions involved from planting to harvesting. Total emissions of CO₂ for growing a poplar SRF with a cycle of eight years and biennial cutting cycle reach a value of 5,910.4 kg ha⁻¹, equivalent to 738.8 kg ha⁻¹ year⁻¹; whereas, for a triennial cutting cycle in a cycle of eight years, the value reached was 5,564.3 kg ha⁻¹, equivalent to 695.5 kg ha⁻¹ year⁻¹. From these results it can be said that growing poplar biomass with triennial cutting cycles involves lower CO₂ emissions compared with growing biomass with biennial cutting cycles.

Considering the yield and the potential energy generated by the biomass, it can be said that the energy produced is 224.1 MJ kg⁻¹ of CO₂ and 297.6 MJ kg⁻¹ of CO₂, respectively for T2 and T3, being T3 the most energy productive cutting cycle.

DISCUSSION

This long-term trial revealed that different harvesting cycles can affect parameters such as biomass yield and carbon balance. The results in this study demonstrate to be consistent with most of the available data, which indicate a strong relationship between the cutting cycle and the productivity of the stand (Deckmyn *et al.* 2004). In fact, for biannual cutting cycles Rafaschieri *et al.* (1999) obtained variable yields between 16 and 20 Mg ha⁻¹ year⁻¹ while for short cutting cycles varying between three and five years, Kauter *et al.* (2003) using *Populus* sp., obtained between 10 and 12 Mg ha⁻¹ year⁻¹ yield.




Figure 2. *Populus deltoides* clone Lux, biomass production during the period 2000-2008.

Producción de biomasa en el cultivo de *Populus deltoides* clon Lux en el periodo 2000-2008.

In our case, CO₂ emissions resulting from the implementation of the various farming operations are minor and contribute in reducing the greenhouse effect caused by the production of energy with other non-renewable fossil fuels. The results of this study can be compared with those obtained by Dornburg *et al.* (2005) for a poplar SRF held in Poland and the Netherlands with fertilization every four years, obtaining values of CO₂ emissions in a range of 1,170 - 1,660 kg ha⁻¹ year⁻¹. In another study, Bacenetti and Fiala (2011), in a poplar SRF in northern Italy with two-year cutting cycle and 18 Mg ha⁻¹ year⁻¹ yields, calculated emissions equivalent to 1,179 kg ha⁻¹ year⁻¹ of CO₂; which is consistent with those results obtained in this study.

From the analyses of the different cutting cycles for a poplar SRF emerges that taking longer cutting cycles consents lower emissions compared with shorter cycles. Comparing these results with others reported by Dubuisson and Sintzoff (1998), it can be shown that depending on the level of intensification, values regarding CO₂ emissions vary from 101-160 MJ kg⁻¹ of CO₂. Whereas the results reported by Matthews (2001) showed mean values of 209.8 MJ kg⁻¹ of CO₂. In the latter study, the calculation of overall energy and carbon budgets required a set of 'standard' assumptions about practices and resultant energy inputs to be made. These standard assumptions accounted for all activities involved in production and delivery of biofuel within the immediate vicinity of the farm (3.2 km). The carbon emissions coefficient exhibited similar sensitivity to input assumptions (Matthews 2001).

Depending on the limits analyzed in the cropping system, emissions will be higher or lower. What is clear is that there is a significant reduction in emissions when producing the same amount of energy with a renewable energy

Table 3. Harvest number, average dry yield (\pm SD), CO₂ emissions and energy produced by poplar SRF for each harvesting cycle (T2 and T3, biennial and triennial cutting cycle, respectively). In the same column different letters stand for significant differences ($P \leq 0.05$).

Número de cosechas, producción media de biomasa seca (\pm SD), emisiones de CO₂ y energía producida para el cultivo de álamo en cada ciclo de corta (T2 y T3, corta bienal y trienal respectivamente). En la misma columna, letras diferentes implican diferencias significativas ($P \leq 0,05$).

Treatment	Parameters			
	Harvest number	Average dry yield (Mg ha ⁻¹ year ⁻¹)	CO ₂ emissions (kg ha ⁻¹ year ⁻¹)	Energy produced (MJ ha ⁻¹ year ⁻¹)
T2	3	9.2 \pm 0.9 b	738.8	165.6
T3	4	11.5 \pm 1.5 a	695.5	207.0

source instead of fossil energy source. To quantify the benefits of renewable energy biomass derived from a poplar SRF with biennial and triennial cutting cycles, we have compared the amounts of CO₂ emitted by the system under study with the amounts released with the use of fossil fuels (table 4). Analyzing table 4, it can be seen how the energy produced per kilogram of CO₂ emitted is much higher in poplar biomass production as an energy source than in the use of any other source of fossil energy. Energy production from biomass is more sustainable, in terms of CO₂ emissions, than the use of fossil energy. These results are similar to those obtained by Sevigne *et al.* (2015) in poplar short rotation forestry in Girona (Spain).

CONCLUSIONS

The poplar SRC is one of the most promising woody species that can be grown in a Mediterranean climate. The study shows that the energy production obtained is higher than the energy needed for the whole productive process,

considering that the clone used is not specific for biomass production. On the other hand, generated CO₂ emissions are low, especially when compared to the emissions generated to produce energy from other fossils energy sources, reducing emissions almost five times. Furthermore, the use of renewable sources of energy contributes in mitigating the CO₂ emissions. Another essential point result suggests that producing energy from poplar biomass can be a sustainable alternative in terms of carbon balance.

As it can be seen, it can be concluded that the use of biomass makes a positive contribution to the reduction of greenhouse gases, in particular CO₂. Finally, the methodology used for calculating the carbon balance can be very useful if applied to any other energy crop.

ACKNOWLEDGEMENTS

We are indebted with Jorge Cuevas Aburto, English coordinator, and Felix W. Doughty, English professor, both from Universidad Santo Tomás, for their valuable corrections of English.

REFERENCES

Table 4. Energy (MJ) produced per kilogram of CO₂ emitted from different energy sources and for a poplar SRF with biennial (T2) and triennial (T3) cutting cycles.

Energía producida por cada kilogramo de CO₂ emitido por diferentes fuentes de energía y para el cultivo de álamo con fines energéticos con turnos de corta bienal (T2) y trienal (T3).

Energy source	MJ produced for CO ₂ kg emitted	Reference
T2	224.1	This study
T3	297.6	This study
Coal	73.3	U.S. Department of Energy 2013
Gasoline motor	54.5	U.S. Department of Energy 2013
Natural gas	72.9	U.S. Department of Energy 2013
Kerosene	53.5	U.S. Department of Energy 2013

- Bacenetti J, M Fiala. 2011. Short Rotation Coppice in Italy: a model to assess economic, energetic and environmental performances of different crop systems. *In* Proceedings of World Renewable Energy Congress. Linköping, Sweden, 8-13 May 2011.
- Bonari E, N Silvestri, S Pampana. 1999. Sisco: a practical tool to assess alternative cropping systems' performances. *In* Proceedings of International Symposium Modelling Cropping Systems. 1999 June 21-23. Lleida, Spain. p. 285-286.
- Cabrera A. 2009. Confronto fra colture "dedicate" per la produzione di biomassa a destinazione energetica in ambiente mediterraneo. PhD Thesis. Pisa, Italy. LAND LAB Scuola Superiore Sant'Anna - Università di Pisa. 173 p.
- Cabrera A, C Tozzini, S Espinoza, R Santelices, E Bonari. 2014. Cálculo del balance energético de una plantación de *Populus deltoides* clon Lux con fines energéticos en un sitio con ambiente mediterráneo. *Bosque* 35(2): 133-139.
- Cannell MGR. 2002. Carbon sequestration and biomass energy offset: Theoretical, potential and achievable capacities globally, in Europe and the UK. *Biomass and Bioenergy* 24 (2): 97-116.

- Deckmyn G, I Laureysens, J Garcia, B Muys, R Ceulemans. 2004. Poplar growth and yield in short rotation coppice: model simulations using the process model SECRETS. *Biomass and Bioenergy* 26: 221-227.
- Dornburg V, G Termeer, APC Faaij. 2005. Economic and greenhouse gas emission analysis of bioenergy production using multi-product crops-case studies for the Netherlands and Poland. *Biomass and Bioenergy* 28: 454-474.
- Dubuisson X, I Sintzoff. 1998. Energy and CO₂ balances in different power generation routes using wood fuel from short rotation coppice. *Biomass and Bioenergy* 15: 379-390.
- ENEA (Agenzia Nazionale per le Nuove Tecnologie, L'energia e lo Sviluppo Economico Sostenibile, IT). 1999. Libro Bianco per la valorizzazione energetica delle fonti rinnovabili. Consulted Jul. 2, 2013. Available in <http://www.regione.sicilia.it/industria/use/Documenti%20ufficiali%20energia/Nazionali/Libro%20bianco%20fonti%20rinnovabili%20Italia%20Aprile%201999.pdf>
- Hoosbeek M, Y Li, G Scarascia-Mugnozza. 2006. Free atmospheric CO₂ enrichment (FACE) increased labile and total carbon in the mineral soil of a short rotation poplar plantation. *Plant and Soil* 281: 247-254.
- IPCC (Intergovernmental Panel on Climate Change, CH). 2006. Guidelines for National Greenhouse Gas Inventories. Consulted Jul. 2, 2013 Available in <http://www.ipcc-nggip.iges.or.jp/public/2006gl/>
- Kaur B, SR Gupta, G Singh. 2002. Carbon storage and nitrogen cycling in silvopastoral systems on a sodic soil in northwestern India. *Agroforestry Systems* 54: 21-29.
- Kauter D, I Lewandowski, W Claupein. 2003. Quantity and quality of harvestable biomass from *Populus* short rotation coppice for solid fuel use - a review of the physiological basis and management influences. *Biomass and Bioenergy* 24(6): 411-427.
- Lewandowski I, A Kicherer, P Vonier. 1995. CO₂-balance for the cultivation and combustion of *Miscanthus*. *Biomass and Bioenergy* 8(2): 81-90.
- Mantineo M, GM D'Agosta, V Copani, C Patane, SL Cosentino. 2009. Biomass yield and energy balance of three perennial crops for energy use in the semi-arid Mediterranean environment. *Field Crops Research* 114: 204-213.
- Matthews RW. 2001. Modelling of energy and carbon budgets of wood fuel coppice systems. *Biomass and Bioenergy* 21 (1): 1-19.
- Rafaschieri A, M Rapaccini, G Manfrida. 1999. Life cycle assessment of electricity production from poplar energy crops compared with conventional fossil fuels. *Energy Conversion and Management* 40: 1477-1493.
- Sevigne E, CM Gasol, F Brun, L Rovira, JM Pagés, F Camps, J Rierdevall, X Gabarrell. 2011. Water and energy consumption of *Populus* spp. bioenergy systems: A case study in Southern Europe. *Renewable and Sustainable Energy Reviews* 15: 1133-1140.
- U.S. Department of Energy. 2013. Independent statistic and analysis. Consulted Jul.3, 2013. Available in <http://www.eia.gov/tools/faqs/faq.cfm?id=73&t=11>.
- West TO, G Marland. 2002. A synthesis of carbon sequestration, carbon emissions, and net carbon flux in agriculture: Comparing tillage practices in the United States. *Agriculture, Ecosystems and Environment* 91(1-3): 217-232.

Recibido: 25.05.15
Aceptado: 29.09.15